

Prenatal Development Timeline

■ Nervous	■ Cardiovascular	■ Muscular	■ Early Events
■ Special Senses	■ Respiratory	■ Skeletal	■ Growth Parameters
■ Blood & Immune	■ Gastrointestinal	■ Endocrine	■ General
■ Skin/Integument	■ Renal/Urinary	■ Reproductive	■ Movement










































Unit 1: The First Week			
Day 0	½ ½	Embryonic period begins	
		Fertilization resulting in zygote formation	
Day 1	½ ½	Embryo is spherically shaped and called a morula comprised of 12 to 16 blastomeres	
		Embryo is spherically shaped with 12 to 16 cells	
Day 1 - Day 1	½ ½	Fertilization - development begins with a single-cell embryo!!!	
Day 2	½ ½	Early pregnancy factor (EPF)	
		Activation of the genome	
		Blastomeres begin rapidly dividing	
		Zygote divides into two blastomeres (24 - 30 hours from start of fertilization)	
Day 3	½ ½	Compaction	
Day 4	½ ½	Embryonic disc	
		Free floating blastocyst	
		Hypoblast & epiblast	
		Inner cell mass	
		See where the back and chest will be	
Day 5	½ ½	Hatching blastocyst	
Day 6	½ ½	Embryo attaches to wall of uterus	
		Solid syncytiotrophoblast & cytotrophoblast	
1 week	½ ½	Chorion	
		Chorionic cavity	
		Extra-embryonic mesoderm (or mesoblast)	
		Placenta begins to form	












Unit 2: 1 to 2 Weeks

1 week, 1 day $\frac{1}{2}$ $\frac{1}{2}$	<ul style="list-style-type: none"> Amnioblasts present; amnion and amniotic cavity formation begins
	<ul style="list-style-type: none"> Bilaminar embryonic disc
	<ul style="list-style-type: none"> Positive pregnancy test
1 week, 2 days $\frac{1}{2}$ $\frac{1}{2}$	<ul style="list-style-type: none"> Corpus luteum of pregnancy
	<ul style="list-style-type: none"> Cells in womb engorged with nutrients
	<ul style="list-style-type: none"> Exocoelomic membrane
	<ul style="list-style-type: none"> Isolated trophoblastic lacunae
	<ul style="list-style-type: none"> Embryonic disc 0.1 mm diameter
1 week, 4 days $\frac{1}{2}$ $\frac{1}{2}$	<ul style="list-style-type: none"> Intercommunicating lacunae network
	<ul style="list-style-type: none"> Longitudinal axis
	<ul style="list-style-type: none"> Prechordal plate
	<ul style="list-style-type: none"> Trophoblastic vascular circle
1 week, 5 days $\frac{1}{2}$ $\frac{1}{2}$	<ul style="list-style-type: none"> Implantation complete
	<ul style="list-style-type: none"> Yolk sac
	<ul style="list-style-type: none"> Embryonic disc diameter: 0.15 to 0.20 mm
1 week, 6 days $\frac{1}{2}$ $\frac{1}{2}$	<ul style="list-style-type: none"> Blood islands in umbilical vesicle
	<ul style="list-style-type: none"> Angiogenesis in chorionic mesoblast
	<ul style="list-style-type: none"> Blood vessels in villi
	<ul style="list-style-type: none"> Connecting stalk
	<ul style="list-style-type: none"> Primordial blood vessels
	<ul style="list-style-type: none"> Amnion with single cell layer
	<ul style="list-style-type: none"> Chorionic villi
2 weeks $\frac{1}{2}$ $\frac{1}{2}$	<ul style="list-style-type: none"> Embryonic epiblast gives rise to primitive streak and primitive node and
	<ul style="list-style-type: none"> Yolk sac
	<ul style="list-style-type: none"> Yolk sac

Unit 3: 2 to 3 Weeks


2 weeks, 1 day $\frac{1}{2}$ $\frac{1}{2}$	<ul style="list-style-type: none"> 3 germ layers
	<ul style="list-style-type: none"> Cloacal membrane
	<ul style="list-style-type: none"> Primitive groove
	<ul style="list-style-type: none"> Rostral-caudal orientation
2 weeks, 2 days $\frac{1}{2}$ $\frac{1}{2}$	<ul style="list-style-type: none"> Erythroblasts in yolk sac

	 Three types of blood-forming cells in yolk sac
	 Primordial germ cells
	 Allantoic diverticulum
	 Allantoic diverticulum
	 Amnion with two cell layers
	 Notochordal process
	 Secondary villi
2 weeks, 4 days  $\frac{1}{2} \frac{1}{2}$	 Foregut, midgut, and hindgut
	 Uteroplacental circulation well established
	 Prechordal plate with 1 retinal field
	 Brain is first organ to appear
	 Caudal eminence
	 Neural ectoderm
	 Neural groove and neural folds
	 Neural plate induced by notochordal process
	 Notochordal and neurenteric canals
	 Notochordal plate
	 Connecting stalk
	 Primitive pit (or notochordal pit)
2 weeks, 5 days  $\frac{1}{2} \frac{1}{2}$	 Prechordal plate with 2 retinal fields
2 weeks, 6 days  $\frac{1}{2} \frac{1}{2}$	 Numerous blood islands in umbilical vesicle
	 Septum transversum (primitive diaphragm)
	 Foregut
	 Oropharyngeal membrane
	 Pharyngeal pouch 1
	 Stomodeum forming
	 Beginnings of the heart can be seen
	 Blood vessels emerge simultaneously in umbilical vesicle, embryo proper, amnion, and connecting stalk
	 Common umbilical artery
	 Dorsal aortae (paired)
	 First pair of aortic arches
	 Heart: Cardiogenic plate, cardiac jelly, myocardial mantle, and endocardial plexus
	 Left ventricle, right ventricle, conotruncus
	 Paired pericardial cavities
	 Paired tubular heart
	 Forebrain, midbrain, and hindbrain
	 Hindbrain with four rhombomeres
	 Isthmus rhombencephali demarcates midbrain and hindbrain


















	 Mesencephalon (or midbrain)
	 Neural cord within caudal eminence
	 Neural groove deepens substantially
	 Primary neuromeres
	 Three main divisions of brain
	 Cephalic and caudal folds
	 Neural crest: Rostral and facial
	 Primitive streak reaches neurenteric canal
	 Somites with central somitocoels: Pairs 1 through 3
3 weeks 	 Blood and blood vessels
$\frac{1}{2}$ $\frac{1}{2}$	








































Unit 4: 3 to 4 Weeks






































3 weeks, 1 day 









































 Thyroid primordium emerges from floor of pharynx

$\frac{1}{2}$ $\frac{1}{2}$

































	 Nephrogenic cord emerges (at 10 somites)
	 Cloaca
	 Common coelomic cavity divides into peritoneal, pericardial, and pleural cavities
	 Liver: Hepatic plate (endoderm)
	 Midgut emerging
	 Pharyngeal arches 1 and 2
	 Pharyngeal cleft 1
	 Second pharyngeal cleft and pouch
	 Pharyngeal groove and ridge with laryngotracheal sulcus
	 Respiratory outgrowth
	 Atria (right and left) far apart
	 Bulbis cordis
	 Circulatory system function begins
	 Endocardial tubes fuse forming tubular heart
	 Heart begins beating
	 Pericardial sac
	 Pericardium
	 Primary head vein
	 Sinus venosus
	 Tubular heart begins folding
	 Umbilical arteries
	 Umbilical veins (right and left)
	 Optic primordia fill neuromere D2
	 Otic pits
	 Chiasmatic plate
	 Mesencephalic flexure
	 Neural tube
	 Neuromeres D1 and D2 (in diencephalon)

	 Optic sulcus in forebrain
	 Pontine region identifiable near cranial nerves VII and VIII
	 Segment D in rhombencephalon
	 Some secondary neuromeres
	 Superior colliculus
	 Telencephalon
	 Telencephalon (or telencephalic) medium
	 Body cavities
	 Hyoid arch
	 Mandibular arch and maxillary process
	 Neural crest: Trigeminal, facioacoustic, glossopharyngeal-vagal, and occipitospinal
	 Somites: Pairs 4 through 12
3 weeks, 3 days $\frac{1}{2}$	 Primordial germ cells begin moving from umbilical vesicle to hindgut
	 Thyroid complete
	 Face: Maxillary and mandibular processes (bilaterally)
	 Cloacal membrane
	 Mesonephric duct emerges from nephrogenic cord
	 Nephric vesicles
	 Cystic primordium
	 Hepatic diverticulum
	 Liver
	 Membrane between future mouth and throat may begin to rupture
	 Angiogenesis along surface of central nervous system
	 Aortic sac
	 Atrioventricular canal
	 Capillary plexus begins forming around brain and spinal cord
	 Conotruncus
	 Conus cordis emerging from right ventricle
	 Endocardium
	 Heart contractions produce peristaltic blood flow
	 Internal carotid arteries
	 Interventricular septum
	 Primordium of myocardium
	 Sinus venosus separating from left atria
	 Trabeculated outpouches along primary cardiac tube representing primordia of left and right ventricles
	 Trigeminal and otic arteries
	 Facio-vestibulocochlear ganglia (CN VII, CN VIII)
	 Glossopharyngeal and vagal ganglia
	 Optic evagination (starting at 14 somites)


	 Otic vesicle
	 Trigeminal ganglia (CN V)
	 Neural crest: Optic crest emerges during Carnegie Stages 11 and 12
	 Nose: Nasal plate
	 Optic vesicles form (17 to 19 somites)
	 Adenohypophysial pouch
	 Adenohypophysis
	 Lamina terminalis
	 Mesencephalon contains tectum and tegmentum
	 Neural crest production and migration continue
	 Neurohypophysial primordia
	 Neuropore (near brain) closes
	 Notochord
	 Segmentation of mesoblast alongside neural tube bilaterally
	 Somites: Pairs 13 through 20
3 weeks, 3 days - 5 weeks, 6 1/2 days	 All eight rhombomeres (Rh 1 through Rh 7, Rh D) - Present in stages 11 through 17
3 weeks, 5 days 1/2	 Telopharyngeal bodies
	 Alimentary epithelium invades stroma of liver
	 Alimentary epithelium proliferates in primordia of stomach, liver, and dorsal pancreas
	 First part of pancreas
	 Gastric portion of foregut elongates (25 to 28 somites)
	 Hepatic primordium with abundant vascular plexus
	 Omental bursa
	 Oropharyngeal membrane is ruptured
	 Pharyngeal arch 3
	 Pharyngeal arches with dorsal and ventral parts
	 Umbilical vesicle elongates
	 Cervical sinus
	 Laryngotracheal groove
	 Lung bud
	 Tracheo-esophageal septum
	 Atrioventricular canal
	 Common cardinal veins (right and left)
	 Descending aorta
	 Heart circulates blood to and from central nervous system, umbilical vesicle, and chorion
	 Hepatocardiac channels (right and left)
	 Rostral and caudal cardinal veins along brain and spinal cord feeding common cardinal veins


	 Septum primum and foramen primum sometimes present
	 Septum primum, foramen primum
	 Sinu-atrial foramen prevents backflow into sinus venosus
	 Sinus venosus collects venous blood from entire embryo
	 Superior vena cava, inferior vena cava, and sinus venosus collecting all venous blood
	 Unidirectional circulation
	 Vitelline arteries and veins
	 Hypoglossal cord (CN XII) enters pharyngeal arch 4
	 Otocyst nearly closed
	 Nasal discs form part of ectodermal ring
	 Optic vesicles covered by sheath (formed by mesencephalic and optic crest)
	 Brain involves 40% of neural tube
	 Brain: Embryonic commissural plate
	 Ectodermal ring complete
	 Hypoglossal nucleus (CN XII)
	 Lowermost spinal cord formation begins
	 Mamillary recess
	 Marginal layer in rhombencephalon
	 Mesencephalic flexure at 90 degrees
	 Mesencephalon with two neuromeres: M1 and M2
	 Motor neurons in basal plate of rhombencephalon
	 Neural tube closes (lower back)
	 Neurofibrils form in rhombencephalon
	 Primary neurulation ends
	 Primordia of ventral thalamus and subthalamus in diencephalon
	 Sulcus limitans
	 Sulcus limitans in midbrain
	 Somites: Pairs 21 through 29
	 Upper limb primordium at level of somites 8 to 10
	 Progressively C-shaped embryo
4 weeks  $\frac{1}{2}$ $\frac{1}{2}$	 Spleen primordia
	 Thymic primordia
	 Lower lip forms from merging of mandibular processes
	 Melanoblasts in epidermis
	 Skin is so thin, you can see through it!
	 Gonadal ridge extends from C-7 to T-8 levels
	 Primordial germ cells migrate to mesonephric ridges
	 Primordial germ cells number several hundred
	 Urorectal septum


	Thyroid bilobed and attached to pharynx by thyroglossal duct
	Diaphragm primordia
	Glomeruli emerge in mesonephros
	Mesonephric duct attached to cloaca
	Nephric tubules now S-shaped
	Urogenital sinus
	Urorectal cleavage line
	Diverticulum ilei marks division between foregut and hindgut
	Esophagus primordia
	Intestines growing in length
	Mesentery from end of duodenum to proximal half of colon
	Opening between gut and umbilical vesicle decreases
	Pancreas: Ventral pancreas
	Pharyngeal pouches 1 through 4
	Pharynx
	Pleuroperitoneal canals
	Small & large intestines
	Stalk of umbilical vesicle lengthens and narrows
	Stomach assumes shape of a spindle
	Umbilical vesicle at height of development
	Vitelline duct
	Bronchial buds
	Lungs begin filling chest cavity
	Mesenchyme from coelomic epithelium surrounds esophagus and lung buds
	Trachea
	Anterior, middle, and posterior cerebral plexuses
	Aorta branches include dorsal intersegmental, lateral segmental, and ventral segmental arteries
	Aortic arches 4 and 6
	Artery from the common iliac artery feeds each lower limb bud
	Atrioventricular bundle
	Cardiac contractions still under myogenic control
	Celiac artery, superior and inferior mesenteric arteries
	Circulatory system "well established"
	Common iliac arteries (right and left, from dorsal aorta bifurcation)
	Contractions well coordinated and sequential from sinus venosus to atria to ventricles
	Functioning two-chamber heart
	Gas exchange through placenta begins
	Gelatinous reticulum (or cardiac mesenchyme)
	Heart chambers bulging with fluid

	 Heart now functions as two parallel pumps
	 Heart rate (about) 113 beats/min
	 Heart: Atrioventricular cushions (rostroventral and caudodorsal)
	 Heart: Myocardium wall 3 to 4 cells thick
	 Primary head veins (right and left) drain anterior, middle, and posterior cerebral plexuses and feed precardinal veins
	 Small arteries emerging throughout mesoderm
	 Ventricle walls trabeculated
	 Vertebral arteries
	 Vitelline veins empty exclusively into hepatic plexus
	 Most cranial nerve ganglia
	 Trigeminal, glossopharyngeal, and vagal preganglia
	 Basement membrane of otic disc surrounds otic vesicle
	 Endolymphatic appendage
	 Otic invagination
	 Otic vesicle closes
	 Terminal-vomerolateral neural crest
	 Brain: Commissural plate
	 Cerebellum
	 Common afferent tract
	 Fourth ventricle
	 Interstitial nucleus (part of medial longitudinal fasciculus)
	 Isthmus rhombencephali (a new neuromere)
	 Oculomotor (CN III) and trochlear nuclei (CN IV) in mesencephalon (midbrain) and isthmus respectively
	 Retinal and lens discs
	 Amnion surrounds connecting stalk and vitelline stalk
	 Amnion surrounds embryo
	 Cervical flexure
	 Hyoid arch subdivides into dorsal and ventral segments
	 Limb buds - the first sign of arms and legs
	 Lower limb buds
	 Umbilical cord emerging
	 Upper and lower limb buds









































Unit 5: 4 to 5 Weeks

4 weeks, 3 days  Early eyes
 $\frac{1}{2} \frac{1}{2}$

4 weeks, 3 days - 5 weeks  Germ cells migrate to gonads
 $\frac{1}{2} \frac{1}{2}$


























4 weeks, 4 days  Thymus
 $\frac{1}{2} \frac{1}{2}$

	Parathyrogenic zones
	Thyroglossal duct
	Thyroid pedicle lengthens
	Dorsal contour develops depression at level of sclerotomes 4 and 5
	Muscular plates between upper and lower limb buds
	Glomerular capsules, partially vascularized
	Mesonephric corpuscle
	Metanephrogenic cap emerges from ureteric bud
	Ureteric buds
	Angiogenesis within peri-esophageal mesenchyme
	Epiploic foramen
	Lesser sac (omental bursa)
	Small intestine forming coils
	Tongue: Hypopharyngeal eminence
	Arytenoid swellings (right and left)
	Capillary network surrounds pulmonary mesenchyme
	Epithelial lamina of larynx
	Lungs: Right and left primary (or main stem) bronchi
	Mesenchyme covering esophagus and respiratory tree separates
	Mesenchyme surrounds bronchi
	Pleura (mesothelium) surrounds part of mesenchyme
	Right main bronchus longer than left
	Atria walls thin, ventricle walls thick and trabeculated
	Atrioventricular cushions not fused
	Common pulmonary vein drains pulmonary plexuses into left atrium
	Conotruncal ridges or cushions (remnants of cardiac jelly)
	Epicardium
	Left subclavian artery feeds left axillary artery, left vertebral artery, and left thyrocervical trunk
	Outflow tract still with one lumen
	Posterior communicating arteries
	Pulmonary arch (sixth aortic arch) forms from aorta and aortic sac
	Pulmonary capillary network fed by pulmonary arteries, drain into left atrium
	Sino-atrial (SA) node
	Superior mesenteric artery and vein
	Upper limb buds with early marginal blood vessel
	Brachial plexus
	Cervical plexus
	Dorsal roots
	Hypoglossal nerve roots unite (CN XII)



















	 Lens and retina invaginate to form optic cup
	 Primordium of cochlear duct
	 Rami communicantes
	 Spinal nerves reach muscle primordia
	 Upper limb buds innervated
	 External ear: Auricular hillocks merging
	 Eyes located on sides of head
	 Lens pits
	 Lens vesicle open to surface (lens pore)
	 Nose: Nasal pits
	 Nose: Nasal plate (or disc) flat or concave
	 Pigment in retina (external layer of optic cup)
	 D1 and D2 no longer identifiable within diencephalon
	 75% of midbrain covered by marginal layer
	 All 16 secondary neuromeres
	 Brain enlarges 50% since Carnegie Stage 13
	 Brain: Cerebral hemispheres appear and begin rapid growth
	 Brain: Lateral ventricles
	 Cerebellum with intermediate and ventricular layers
	 Cerebellum: Primordium found in alar plate of rhombomere 1
	 Corpora striata primordia connected by commissural plate
	 Cranial nerve 3
	 Di-telencephalic sulcus
	 Dorsal and ventral thalami
	 Dorsal funiculus
	 Hypothalamic sulcus
	 Hypothalamus
	 Mamillary region
	 Medial and lateral longitudinal fasciculi
	 Median ventricular eminence
	 Pontine flexure
	 Preoptic sulcus extends between optic evaginations
	 Preoptico-hypothalamo-tegmental tract
	 Primary meninx surrounds most of brain
	 Rhombic lip
	 Spinal cord wall with three zones: ventricular (ependymal) zone, mantle (intermediate) zone, and marginal zone
	 Subthalamus with medial striatal ridge emerging
	 Synencephalon
	 Tegmentum
	 Tentorium cerebelli, medial portion






































	■ Terminal-vomeronasal crest contacts brain (olfactory area)
	■ Torus hemisphericus (TH)
	■ Velum transversum
	■ Ventral longitudinal fasciculus
	■ Ventral segment of hyoid arch subdivides
4 weeks, 5 days $\frac{1}{2}$ $\frac{1}{2}$	■ Primordium of antitragus emerges from ventral subsegment of hyoid arch
	■ Gonad framework found in coelomic epithelium
	■ Thyroid detached from epithelium of pharynx in some embryos
	■ Lower limb bud rounded proximally and tapered distally
	■ Mesenchymal skeleton in upper and lower limbs
	■ Right and left neural processes
	■ Sclerotomic material around notochord (rhombomere D level)
	■ Vertebrae well defined
	■ Vertebral centra
	■ Primary urogenital sinus
	■ Ureteric bud extends to pelvis of the ureter
	■ Bladder and rectum are separating caudal to ureters
	■ Caecum
	■ Dense mesenchyme surrounds much of gastrointestinal tract
	■ Esophagus elongates, passes dorsal to carina and between main stem bronchi
	■ Gall bladder and cystic duct
	■ Liver: Hepatic ducts
	■ Ventral pancreas appears as an offshoot of the cystic duct
	■ Lobar bud swellings denote areas of secondary bronchi
	■ Remnants of coelomic epithelium forming visceral pleura
	■ Atrioventricular cushions apposed
	■ Blood flow divided into right and left streams through atrioventricular canal, ventricles, outflow tract, and aortic sac
	■ Blood vessels penetrate diencephalon
	■ Capillary plexus surrounds esophagus
	■ Capillary plexus surrounds lung buds
	■ Cardiac mesenchyme surrounds ventricles and outflow tract
	■ Coronary arteries (terminal end)
	■ Foramen secundum begins in septum primum
	■ Left ventricle with thicker walls and greater volume than right

	<div></div> Right subclavian artery originates from brachiocephalic artery and feeds right thyrocervical trunk and axillary and vertebral arteries
	<div></div> Semilunar cusps
	<div></div> Capsule present around lens
	<div></div> Corneal epithelium overlying optic cup
	<div></div> Ear: Endolymphatic duct
	<div></div> Geniculate and vestibulocochlear ganglia separating
	<div></div> Lens body now present containing some lens fibers
	<div></div> Lower limb buds innervated
	<div></div> Optic stalk
	<div></div> Utricle, endolymphatic duct, and endolymphatic sac
	<div></div> Utriculo-endolymphatic fold
	<div></div> External ear primordia emerges from caudolateral portion of mandibular arch
	<div></div> Face: Lateral and medial nasal processes bilaterally
	<div></div> Lateral nasal processes along dorsolateral lip of nasal pits
	<div></div> Lens vesicles closed, pores absent
	<div></div> Nose: Nasal discs recede forming nasal pits
	<div></div> Optic chiasm
	<div></div> Adult lamina terminalis
	<div></div> Amygdaloid area
	<div></div> Brain with five main sections
	<div></div> Cerebellar plate
	<div></div> Cerebellum with marginal layer
	<div></div> Fibers of dorsal funiculus reach level of C1
	<div></div> First axodendritic synapses in cervical spinal cord
	<div></div> First nerve fibers
	<div></div> Habenular nucleus
	<div></div> Habenulo-interpeduncular tract
	<div></div> Lateral striatal ridge (derived from telencephalon and comprised mainly of neostriatum)
	<div></div> Lateral ventricular eminence
	<div></div> Locus caeruleus
	<div></div> Longitudinal zones in diencephalon
	<div></div> Marginal layer throughout most of diencephalon
	<div></div> Material for sympathetic trunks scattered in cervical region
	<div></div> Median striatal ridge (paleostriatum)
	<div></div> Mesencephalic tract of CN 5
	<div></div> Most cranial nerves seen
	<div></div> Olfactory fibers reach brain
	<div></div> Optic groove (also called preoptic recess)
	<div></div> Postoptic recess
	<div></div> Primordium of epiphysis



































	 Rhombomeres still identifiable
	 Superior colliculi and its commissure
	 Superior medullary velum
	 Supramamillary commissure
	 Synapses among motor neurons in spinal cord
	 Tectobulbar tract
	 Tentorium
	 Third ventricle
	 Trigemino-cerebellar tract
	 Trochlear nerve root and decussation (CN IV)
	 Hand plate emerges from distal upper limb bud
	 Frontonasal prominence
5 weeks  	 ACTH [adrenocorticotropin hormone]
$\frac{1}{2}\frac{1}{2}$	
	 Growth hormone
	 Pituitary gland
	 Limb buds form hand plates
	 Permanent kidneys
	 Arytenoid and epiglottal swellings
	 Bronchial tree branching accelerates
	 Lobar pattern mimics adult pattern
	 T-shaped laryngeal inlet
	 Pacemaker cells
	 Head is one third of entire embryo

Unit 6: 5 to 6 Weeks





























5 weeks, 1 day  	 Wrist joints are forming
$\frac{1}{2}\frac{1}{2}$	
5 weeks, 2 days  	 Apical epidermal ridges
$\frac{1}{2}\frac{1}{2}$	
	 Mammary ridge
	 Maxillary and premaxillary fields still widely separated
	 Nipples emerge from mammary crest
	 Gonad region separates from mesonephros
	 Gonadal primordium
	 Labioscrotal swelling
	 Urogenital fold and groove
	 Suprarenal gland: Cortex primordium
	 Suprarenal gland: Medulla
	 Thyroid detaches from pharynx
	 Thyroid with right and left lobes connected by an isthmus
	 Cartilage in mandibular arch









































	 Hand area with central carpal region and digital plate with marginal vein
	 Pre-chondrocranium: Otic capsule, nasal capsule, and parachordal condensations
	 Primordia of primary palate
	 Ribs: Primordia now present for all 12 pairs
	 Vertebral column with 36 levels of ganglia and myotomes
	 Extra-ocular premuscle masses receive cranial nerve fibers [oculomotor (CN III), trochlear (CN IV), and abducens (CN VI) nerves]
	 Gluteal mesoderm
	 Infrahyoid premuscle masses
	 Limb mesoderm
	 Sternocleidomastoid-trapezius premuscle mass with spinal accessory nerve (CN11)
	 Thigh and thigh mesoderm
	 Tongue premuscle mass
	 Metanephros at level of sacrum
	 Urethral plate
	 Lesser omentum (ventral mesogastrum)
	 Peritoneal cavity
	 Rectum
	 Stomach: Greater and lesser curvatures
	 Yolk stalk disappears
	 Bronchial tree expanding
	 Cervical sinus diminished in size
	 Epiglottis
	 Primitive Larynx
	 Anterior, middle, and posterior cerebral arteries
	 Atrioventricular (AV) node
	 Atrioventricular cushions fuse with interventricular septum
	 Circle of Willis almost complete
	 Conotruncal septum
	 Endocardial cushions (rostroventral and caudodorsal) begin fusing around atrioventricular canal forming right and left atrioventricular canals and two separate blood streams
	 External carotid artery
	 Foramen primum disappearing
	 Hepatic portal vein
	 Infundibulum of right ventricle
	 Jugular lymph sac
	 Lateral atrioventricular cushions
	 Mesencephalic artery
	 Myelencephalic artery

	Perilental blood vessels
	Primitive cavernous sinus drains primitive maxillary and supraorbital veins
	Primitive renal plexus
	Right ventricle feeds sixth (pulmonary) aortic arches; left ventricle feeds fourth aortic arches
	Semilunar valves (aortic and pulmonary) are forming
	Ventricles each with three parts: inlet, trabecular pouch, and outflow tract
	Ventricles enlarge and deepen side-by-side forming an ever growing interventricular septum
	Celiac plexus
	Cochlear nerve present
	Femoral and obturator nerves innervate rostrolateral part of lower limb
	Hypoglossal nerve (CN XII) reaches tongue
	Intercostal nerves
	Lumbar and sacral plexuses
	Musculocutaneous, radial, ulna, and median nerves enter upper limb bud
	Nasal pits face more ventrally, still widely separated
	Nasofrontal groove
	Olfactory fibers connect nasal pits with brain
	Olfactory fibers enter brain
	Olfactory tubercle present
	Peroneal and tibial nerves innervate caudomedial part of lower limb
	Phrenic nerve
	Pigment in retina visible externally
	Primordium of cochlear pouch
	Tibial nerve innervates foot area
	Auricular hillocks on hyoid arch (antitragus and helix)
	Auricular hillocks on mandibular arch (tragus and crus)
	Blind nasal sac
	Nasal fin
	Alar lamina emerging with dense rhombic lip
	All cranial nerves identifiable
	Archipallium, paleopallium, and neopallium
	Area epithelialis
	Brain: Primordial plexiform layer in area of future temporal lobe
	Cajal-Retzius cells
	Commissure of the trochlear nerve
	Diencephalic subthalamic nucleus
	Dorsal and ventral thalami separated by groove
	Dorsal funiculus fibers reach medulla oblongata
	Epiphysis cerebri

	 Glial cells identifiable adjacent to neurons
	 Greater petrosal nerve
	 Hippocampus: Gyrus dentatus
	 Infundibular recess and infundibulum
	 Interventricular foramen large
	 Marginal ridge
	 Medial and lateral ridges of corpus striatum are continuous
	 Median forebrain bundle
	 Neurohypophysial outgrowth
	 Olfactory tubercle
	 Pontine flexure deepens
	 Posterior commissure
	 Recurrent laryngeal nerve
	 Reticular formation more defined
	 Retinal fissure closes
	 Splanchnic nerve
	 Sulcus limitans hippocampi
	 Superior laryngeal nerve
	 Second pharyngeal arch more prominent
	 Third pharyngeal arch recedes
5½ weeks ½½	 Initial tooth formation
5½ weeks - 6 weeks ½½	 Subtle movement begins
5 weeks, 4 days ½½	 Cartilage formation
5 weeks, 5 days ½½	 Nerve cells differentiating
5 weeks, 5 days - 7 weeks, 1 day ½½	 Melanocytes in epidermis
5 weeks, 6 days ½½	 Facial growth centers grow and begin merging forming nose and upper jaw
	 Genital eminence forms phallus or genital tubercle
	 Gonad grows into oval shape with irregular surface
	 Auditory ossicles identifiable in mesenchyme
	 Cartilage in occipital sclerotomes (1-4)
	 Digital rays in hand plate
	 Femur: Chondrification begins
	 Foot with rounded digital plate
	 Hypoglossal foramen (or canal) through sclerotome 4 (area of future occipital bone)

	Odontogenic epithelium emerges in six areas (four maxillary and two mandibular)
	Primary palate components (right and left) fuse in midline
	Primitive palatine groove
	Primordium of cartilage within nasal septum
	Vertebral centra begin chondrification
	Primordia of orbital muscles
	Calices
	Mesonephros can produce urine
	Pelvis of the ureter with three main divisions
	Vesico-urethral canal
	Biliary ducts within liver
	Dorsal and ventral pancreas fuse but retain separate ducts
	Duodenum enlarges proximal to and distal to bile and pancreatic ducts
	Esophagus developing a submucous coat surrounding epithelium
	Intestinal loop begins umbilical herniation
	Primordial vermiform appendix
	Stomach regions include gastric canal, fundus, corpus (or body), and pyloric antrum
	Trachea: Precursors of tracheal cartilages
	Condensing mesenchyme around junction between left and right atria and cardiac tube is precursor to mitral and tricuspid valves
	Outflow tract rotates counterclockwise
	Right and left atrioventricular canals totally separated
	All parasympathetic cranial nerve ganglia identifiable
	All spinal nerves present
	Cell islands in olfactory tubercle
	Crescentic lens cavity
	Geniculate ganglion separate from vestibulocochlear nerve
	Globular process emerges from each medial nasal process
	Nasal fin connecting nasal disc and surface epithelium
	Nasofrontal grooves
	Olfactory tubercle with cellular islands
	Hyomandibular groove enlarges (onset of concha and external auditory meatus formation)
	Medial rims of nasal pits form nasal septum
	Nostril becomes continuous with nasal sac
	Primary lens fibers
	Retinal fissure closed
	Capillaries between adenohypophysis and hypothalamus

	 Commissure of the oculomotor nerves
	 Cortical nucleus in amygdaloid body
	 Dentate and isthmic nuclei in cerebellum
	 Dura begins forming in basal area
	 Epiphysis cerebri with intermediate layer
	 First hint of septal nucleus
	 Frontal and temporal poles of cerebral hemispheres
	 Gustatory fibers separate from common afferent tract
	 Hemispheric stalk
	 Intermediate layer in tectum mesencephali
	 Interventricular foramen
	 Mesencephalon with intermediate layer
	 Somites: Pairs 38 and 39
	 Spinal cord reaches caudal tip of body
	 Subarachnoid space
	 Synapses in spinal cord between interneurons and primary afferent neurons
	 Ventral thalamus with intermediate layer
	 Anterior choroid artery
6 weeks  	 Face withdraws from light touch around mouth
	 Blood forming in liver
	 Milk lines
	 Nipples along side of trunk
	 Adrenal glands
	 Glucagon in pancreas
	 Handplates develop subtle flattening
	 Joints
	 Medial skull cartilages: Parachordal, hypophyseal, and trabecular
	 Tooth buds (primary teeth)
	 Diaphragm is largely formed
	 Intestines fill base of umbilical cord
	 External ears
	 Synapses form in spinal cord
	 Crown-heel length 1.6 cm
Unit 7: 6 to 7 Weeks	
6 weeks, 2 days  	 Angiogenesis begins inside gonads
	 Gonad grows into oval shape with irregular surface
	 Ostium (abdominal) of uterine tube at rostral end of paramesonephric duct (in female embryos)
	 Paramesonephric duct forms from rostral end of mesonephric duct

	 Testicular cords in gonads of male embryos
	 Testicular cords in male gonad
	 Elbow regions sometimes identifiable
	 Embryo with cervical and lumbar flexures
	 Embryo with dorsal concavity
	 Finger rays with early interdigital notching
	 Hands polygon-shaped
	 Humerus, radius, and ulna
	 Humerus: Chondrocytes in phases one through three
	 Scapula and clavicle
	 Semicircular ducts form in order: anterior, posterior, and lateral
	 Sternum: Episternal cartilage created from fusion of right and left sternal bars
	 Tibia and fibula
	 Toe rays sometimes present
	 Deltoid muscle
	 External and internal abdominal oblique muscles
	 Levator scapulae muscle
	 Longus cervicis and semispinalis cervicis muscles
	 Pectoralis major muscles
	 Platysma muscle
	 Rectus abdominis muscle
	 Rectus capitis posterior and semispinalis capitis muscles
	 Serratus anterior muscles
	 Splenius and longissimus muscles
	 Stapedius muscle
	 "Common excretory duct is disappearing"
	 Cloacal membrane ruptures (stages 18-19)
	 Primordia of secretory tubules
	 Esophagus with muscular and submucous coats
	 Submandibular gland primordia
	 Bronchial tree with subsegmental buds
	 Bronchial tree with well established segmental bronchi
	 Lingula of left upper lobe
	 Aortic and pulmonary valves assuming shape of a cup
	 Brachiocephalic veins, right and left
	 Inferior vena cava
	 Interventricular septum: membranous part begins forming
	 Left coronary artery arises from aorta
	 Mesenchyme ridges in place of future mitral and tricuspid valves
	 Pulmonary and aortic blood flows completely separate






	Secondary interventricular foramen sometimes closing (stage 18-21) interventricular septum
	Septum secundum and foramen ovale (stages 18-21)
	Bucconasal membrane
	Bucconasal membrane detaches opening up nasal airway
	Crus commune
	Ethmoidal epithelium emerges from upper medial nasal wall
	Frontonasal angle (marks location of future nasal bridge)
	Mesenchyme thickenings mark beginning of "sclera and its muscular attachments"
	Nasal tip emerges
	Nerve fibers in retina
	Optic fibers
	Retina's outer lamina heavily pigmented
	Vomeronasal nerve and ganglion
	Vomeronasal organ marked by groove and located in fold of lower medial nasal wall
	Choanae
	Conjunctival sac marked by groove
	Cornea and conjunctiva
	Ear: Stapes primordium surrounds stapedia artery
	External ear: Crus helices forming from auricular hillocks two and three (from mandibular arch)
	Eyelid folds sometimes present
	Nasal fin splits forming choanae and bucconasal membrane
	Nasolacrimal duct begins as epithelial strand emanating from nasomaxillary groove
	Nostrils, nasal wings, and nasal septum easily seen
	Olfactory bulb sometimes with olfactory ventricle
	Primary lens fibers filling lens vesicle cavity
	Adenohypophysis no longer open to pharyngeal cavity
	Archistriatum
	Brain: Dentate nucleus in internal cerebellar swellings
	Brain: Pineal recess emerges representing anterior lobe of epiphysis
	Brainwave activity has begun
	Cerebrospinal fluid production begins
	Choroid plexuses in fourth and lateral ventricles
	Corpus striatum much larger extending to preoptic sulcus; has subtle groove
	External cerebellar swellings contain future flocculus
	Four amygdaloid nuclei
	Fourth ventricle: Choroid folds





































	Hippocampus reaches olfactory region
	Interpeduncular fossa
	Neurohypophysis walls are folded
	Nucleus ambiguus of the vagus (CN10)
	Prosencephalic septum
	Red nucleus
	Substantia nigra
	Supra-optic commissure
6½ weeks ½ ½	The hands begin to move
	Volar pads on palms
	Bones first form in the collar bones and lower jaw
6 weeks, 5 days ½ ½	Greater thymic bud
	Cheeks form by merging of maxillary and mandibular processes
	Mammary gland primordium
	Mammary ridge disappears leaving only mammary gland primordium
	Female duct
	Gonads extend from levels T-10 to L-2
	Rete ovarii (in female embryos)
	Rete testis begins emerging from seminiferous cords (Stage 19-23) (in male embryos)
	Tunica albuginea in male embryos
	Suprarenal gland: Cortex
	Suprarenal gland: Medulla populated by prechromaffin cells
	Arms point forward
	Beginnings of occipital and sphenoid bones
	Bilateral cartilaginous sternal bars tie ribs together; sternal bars join cranially to form the episternal bar in the midline
	Cartilage within otic capsule envelops semicircular canals and cochlear duct
	Cartilaginous styloid process
	Ear: Cartilaginous malleus, incus, and stapes (the middle ear ossicles)
	Ectomeninx covers lateral and dorsal surfaces of brain (laying the foundation for the flat bones of the skull)
	Intervertebral discs form from caudal condensed portion of sclerotomes
	Ischium and ilium
	Labiodental lamina: Inner dental lamina and outer labi gingival band
	Laryngeal cartilages











































	<input type="checkbox"/> Limbs point forward (ventrally)
	<input type="checkbox"/> Orbitosphenoid cartilage located within ectomeninx near optic stalk
	<input type="checkbox"/> Ossification begins in maxilla (stages 19 -20)
	<input type="checkbox"/> Primitive palate (or intermaxillary segment)
	<input type="checkbox"/> Rib primordia become cartilaginous
	<input type="checkbox"/> Ribs each have an identifiable head and shaft
	<input type="checkbox"/> Trachea: Tracheal cartilage
	<input type="checkbox"/> U-shaped labiodental lamina form along upper and lower oral cavity
	<input type="checkbox"/> Vertebral column represented by cartilaginous centrum, neural arch, and short transverse process
	<input checked="" type="checkbox"/> Esophagus: Muscularis layer adjacent to esophageal plexus
	<input checked="" type="checkbox"/> Gluteal muscle group
	<input checked="" type="checkbox"/> Iliopsoas muscles
	<input checked="" type="checkbox"/> Infrahyoid muscles
	<input checked="" type="checkbox"/> Internal intercostal muscles
	<input checked="" type="checkbox"/> Limb extensor muscles located dorsally
	<input checked="" type="checkbox"/> Limb flexor muscles located ventrally
	<input checked="" type="checkbox"/> Midgut: Muscularis
	<input checked="" type="checkbox"/> Muscle tissue forming around phrenic nerve within septum transversum portion of diaphragm
	<input checked="" type="checkbox"/> Pharyngeal constrictor muscle
	<input checked="" type="checkbox"/> Premuscle mass of the muscles of mastication innervated by mandibular nerve
	<input checked="" type="checkbox"/> Quadratus lumborum muscle
	<input checked="" type="checkbox"/> Rhomboid and scalene muscles
	<input checked="" type="checkbox"/> Sternocleidomastoid and trapezius muscles distinct and innervated by separate branches of spinal accessory nerve (CN XI)
	<input checked="" type="checkbox"/> Thenar and hypothenar eminences
	<input checked="" type="checkbox"/> Tongue forms from swellings in floor of pharynx
	<input checked="" type="checkbox"/> Tongue: Extrinsic muscles identifiable
	<input checked="" type="checkbox"/> Tongue: Intrinsic muscles identifiable
	<input checked="" type="checkbox"/> Transversospinal and erector spinae muscle groups
	<input checked="" type="checkbox"/> Upper limb flexors innervated by musculocutaneous, median, and ulnar nerves
	<input checked="" type="checkbox"/> Major calyces, cranial and caudal, with collecting tubules within metanephrogenic mass
	<input checked="" type="checkbox"/> Mesonephros extends from T-9 to L-3
	<input checked="" type="checkbox"/> Metanephros extends from T-12 to L-2
	<input checked="" type="checkbox"/> Renal capsule covers distal collecting tubules
	<input checked="" type="checkbox"/> Renal vesicles form in part of metanephros
	<input checked="" type="checkbox"/> Ureter forms from "proximal segment of metanephric diverticulum"
	<input checked="" type="checkbox"/> Urogenital sinus comprised of three parts: Bladder, pelvic, and phallic portions









































	■ Anal folds adjacent to anal membrane
	■ Anal membrane
	■ Duodenum: "Assumes the shape of an arc"
	■ Greater omentum
	■ Lateral palatine process
	■ Liver: rapid growth, right side greater than left
	■ Median mandibular groove disappears as mandibular processes merge in midline
	■ Palatine fossa (from pharyngeal pouch 2)
	■ Primitive oral cavity
	■ Primitive rima oris replaces stomodeum
	■ Stomach wall layers: Mucosa, submucosa, muscularis, and serosa
	■ Submandibular and parotid gland buds
	■ Submandibular gland duct
	■ Bronchial tree: First generation of subsegmental bronchi complete
	■ Glottis, primitive
	■ Lung sac, right: Oblique and horizontal fissures define upper, lower, and middle lobes
	■ Lung sac: Apex and base
	■ Lung, left: Oblique fissure defines upper and lower lobes
	■ "Septum primum fuses with endocardial cushions" obliterating ostium primum and creating the ostium secundum
	■ Apex of left ventricle
	■ Circulus arteriosus (Circle of Willis) complete
	■ External iliac arteries
	■ Iliac lymph sac
	■ Intercostal and subcostal arteries
	■ Internal thoracic artery and costocervical trunk
	■ Mesenteric lymph sac
	■ Mesonephric artery feeds mesonephros, gonads, and suprarenal glands
	■ Papillary muscles
	■ Pontine, superior cerebellar, and anterior and posterior inferior cerebellar arteries replace myelencephalic and metencephalic arteries
	■ Primitive marginal sinus drains diencephalon
	■ Primitive tentorial sinus drains cerebral vesical
	■ Primitive transverse and sigmoid sinuses
	■ Pulmonary arteries (right and left)
	■ Right coronary artery arises from aorta
	■ Splenic vein
	■ Tricuspid and mitral valves

	Anterior chamber between iridopupillary membrane and thickened ectoderm
	Auditory tube and primitive tympanic cavity form from tubotympanic recess pharyngeal pouch 1)
	Celiac, superior mesenteric, and inferior mesenteric preaortic ganglia
	Choana
	Cochlear duct tip grows upward
	Esophageal plexus formed by vagal nerves (CN X)
	Facial nerve (CN VII) branches: Chorda tympani, greater petrosal, posterior auricular, and digastric
	Facial nerve (CN VII) reaches cervicomandibular region
	Glossopharyngeal nerve (CN IX) innervates stylopharyngeus premuscle mass
	Hypoglossal nerve (CN XII) innervates separating tongue muscles
	Linguogingival groove
	Nasolacrimal duct forms from maxillonasal groove
	Nasolacrimal ducts extend from medial eyes to primitive nasal cavity
	Nerve fibers begin extending from retina
	Optic fibers enter chiasmatic plate
	Primitive nasal cavity
	Primordial vitreous body
	Superior, middle, and inferior cervical ganglia
	Trigeminal nerve (CN V) with ophthalmic, maxillary, and mandibular divisions reach their destinations
	Vagal trunks, anterior and posterior, extending into abdomen
	Eyelids: Upper and lower lids present and growing
	Saccul and cochlear duct
	Adenohypophysis: Lateral lobes of pars tuberalis
	Adenohypophysis: Pars intermedia emerging
	Brain: Internal capsule formation underway
	Cerebral hemispheres cover half of diencephalon
	Dorsal and ventral cochlear nuclei
	Fourth ventricle: Lateral recesses
	Ganglion of nervus terminalis
	Globus pallidus externus in the diencephalon
	Habenular commissure
	Intermediate layer in dorsal thalamus
	Lemniscal decussation
	Lower limb nerves (femoral, obturator, sciatic, common peroneal, and tibial) identifiable
	Medial accessory olivary nucleus
	Neurohypophyseal bud
	Nuclei of forebrain septum







































	 Nucleus accumbens
	 Occipital pole of cerebral hemispheres
	 Optic stalk with barely discernible lumen
	 Paraphysis marks dividing line in roof between telencephalon and diencephalon
	 Primitive filum terminale
	 Radial nerve innervates upper limb extensors
	 Rhombomeres no longer distinguishable
	 Subcommissural organ
	 Zona limitans intrathalamica between dorsal and ventral thalami
6 weeks, 6 days   	 Feet polygon-shaped
	 Cloacal membrane ruptures
7 weeks  	 Head rotates
	 Leg movements
	 B lymphocytes in liver
	 Ovaries
	 Testes begin to differentiate
	 Insulin in pancreas
	 Foot plates notched
	 Hiccups
	 Tendons attach muscle to bone
	 The heart has four chambers and is nearly complete.
	 The heart rate peaks at 165 to 170 beats per minute.
	 Crown-heel length 2.2 cm
Unit 8: 7 to 8 Weeks	
7 weeks, 1 day  	 Facial processes no longer distinguishable
	 Ovaries full of primitive oogonia, intermediate pregranulosa cells, and mesenchyme
	 Testes with short straight tubules
	 Upper limbs with slightly flexed elbows
	 Diaphragm: Central tendon
	 Renal vesicles with S-shaped lumina
	 Submandibular gland: Solid epithelial ducts enlarge and begin to branch
	 Adenohypophysis with new capillaries on rostral surface
	 Scalp vascular plexus
	 Cochlear duct tip growing horizontally
	 Lens cavity completely filled
	 Optic commissure






























	 Optic fibers extend to optic chiasma
	 Cornea with three layers
	 Brain: Inferior colliculus (in mesencephalon)
	 Cerebral hemispheres expand beyond lamina terminalis
	 Cerebral hemispheres extend over two-thirds of diencephalon
	 Interpeduncular groove
	 Medial septal nucleus
	 Nigrostriatal fibers
	 Nucleus of diagonal band
	 Sacrocaudal spinal cord formation (secondary neurulation) complete
	 Sensory pathways: Cuneate and gracile decussating fibers
	 Septum verum
	 Spinothalamic tract
7 weeks, 1 day - 8 weeks 1/2 1/2	 Stomach: Folds in stomach wall
7 weeks, 2 days 1/2 1/2	 Arteries and veins of heart complete
7 weeks, 3 days 1/2 1/2	 Volar pads begin to emerge on fingertips
	 Chondrocranium with dorsum sellae and hypophysial fossa
	 Dens (of second cervical vertebrae)
	 Sternoclavicular joint and manubrium
	 The knee joints have arrived
	 Trachea: Thyroid cartilage
	 Wrists slightly flexed
	 Gluteus medius and gluteus minimus muscles
	 Iliacus muscles
	 Mylohyoid and infrahyoid muscles
	 Orbicularis oculi muscles
	 Submandibular gland: Solid ducts with definitive branches
	 Anterior and posterior choroid arteries
	 Left superior vena cava disappears (Stages 21-23)
	 Scalp vascular plexus moving toward vertex
	 Cornea: Substantia propria layer
	 Fibers of optic nerve reach brain
	 Eyelids growing rapidly
	 Anterior and inferior horns of lateral ventricle
	 Brain: Insula within cerebral hemisphere
	 C-shaped lateral ventricle

	 Cerebral hemispheres cover 75% of diencephalon
	 Cerebral hemispheres cover more than half of diencephalon
	 Cortical plate within primordial plexiform layer
	 Glial and neurilemmal (Schwann) cells within cranial nerves
	 Globus pallidus internus
	 Internal fiber layer of cerebellum
	 Lateral olfactory tract
	 Primordium of dentate nucleus
	 Pyramidal cells in hippocampus
	 Subthalamic nucleus proper, entopeduncular nucleus, and globus pallidus externus within subthalamus
	 Sulcus transversus rhombencephali
	 Ventral part of lateral geniculate body
7½ weeks  	 Hands begin to touch face
	 The hands touch each other as do the feet!
	 Fingertips thicken
	 Plantar pads toes
	 EKG pattern similar to adult
7 weeks, 4 days  	 The fingers are free
7 weeks, 5 days  	 Bone-forming cells called osteoblasts emerge
	 Bone-forming cells emerge
	 Endolymphatic and jugular foramina
	 Hands can reach one another and fingers can overlap
	 Optic foramen, foramen rotundum, internal acoustic foramen
	 Osteoblasts emerge
	 Pelvis: Obturator foramen
	 Obturator internus muscles
	 Rectus femoris muscle
	 Large glomeruli present within metanephros
	 Submandibular gland: Secondary branching with lumen formation starting at oral end of duct
	 Costodiaphragmatic recess of pleural cavity
	 Chordae tendineae (Stages 22 and 23)
	 Intradural veins (sinuses)
	 Scalp vascular plexus 75% of the way to the vertex
	 Cochlear duct's second loop growing upward
	 Scleral condensation
	 Tragus and antitragus taking shape

	 Eyelids continue growing rapidly over the surface of the cornea
	 Optic nerve acquires a sheath
	 Brain: Claustrum
	 Brain: Cortical plate within cerebral hemispheres
	 Brain: Internal capsule with connections to epithalamus, dorsal thalamus, and mesencephalon
	 Brain: Putamen
	 Cerebral hemispheres cover 75% of diencephalon
	 Commissural plate thickens
	 Cortical plate expanding rapidly
	 Folds in roof of third ventricle
	 Nerve fibers between neopallial subplate and internal capsule
	 Thalamocortical fibers
7 weeks, 6 days   $\frac{1}{2} \frac{1}{2}$	 The toes are free
8 weeks   $\frac{1}{2} \frac{1}{2}$	 Complex response to touch
	 More frequent hand-to-face contact
	 Mouth opens & closes
	 Squinting
	 The embryo floats and rolls over in the womb
	 Hairs first appear in eyebrows & around mouth
	 Skin multi-layered, loses transparency
	 Ductus deferens
	 Interstitial cells forming within testis
	 Testicular tubules
	 Male embryos are making testosterone already!
	 Anterior inferior iliac spine
	 Costal cartilage
	 Enamel organ
	 Femur: Head and acetabular fossa
	 Glenoid fossa
	 Greater trochanter
	 Head of humerus
	 Inguinal ligament
	 Joint development: Cavitation underway in hip, knee, and ankle (in some embryos)
	 Joint development: Cavitation underway in shoulder, elbow, and wrist (in some embryos)
	 Nucleus pulposus (from notochord)
	 Ossification underway in scapula and distal phalanges in some embryos
	 Pubic symphysis

	□ Scapular spine and notch
	□ Skull: Foramen magnum (wide)
	□ Skull: Ossification underway in some embryos
	□ Superior and inferior pubic rami
	□ The embryo's joints are similar to adult joints
	□ Ulna: Styloid process and olecranon
	□ Vertebrae cartilaginous (33 or 34 in number)
	■ Anterior digastric muscles
	■ Depressor anguli oris muscle
	■ Diaphragm complete
	■ Esophagus: Longitudinal muscles
	■ Obliquus superior capitis muscle
	■ Obturator externus, gluteus maximus, and hamstring muscles
	■ Posterior belly of the digastric muscle
	■ Psoas tendon
	■ Rectus sheath with anterior and posterior lamina
	■ Temporal and lateral pterygoid muscles
	■ Zygomaticus major muscle
	■ Kidneys at level of first three lumbar vertebrae
	■ Metanephros: Numerous large glomeruli
	■ Metanephros: Secretory tubules elongating and becoming convoluted
	■ Sinus tubercle
	■ Urethra
	■ Urine production and release
	■ Gastrosplenic ligament
	■ Nerves reaching intestinal loop
	■ Peristalsis in large intestine
	■ Submandibular gland: Lumen present in terminal portions of duct
	■ Submandibular gland: Mesodermal sheath surrounds gland
	■ Unfused uvula (edge of unfused palatine shelf) and secondary palate
	■ Occasional breathing motions begin
	■ Pseudoglandular stage begins
	■ Azygos vein
	■ Blood supply to the brain closely resembles adult pattern
	■ Hemiazygos veins
	■ Inferior epigastric artery
	■ Inferior vena cava valve at junction of right atrium
	■ Scalp vascular plexus nearing vertex
	■ Submandibular glands: Angiogenesis begins around epithelial tree (ducts)


























	 Superior sagittal sinus
	 Cochlear duct's 2.5 coils nearly complete
	 Cranial nerve distribution mimics adult pattern
	 Ear drum
	 Eye: Secondary vitreous body
	 Lens: Secondary lens fibers emerging
	 Retina: Eight layers present
	 Retina: Four of the ten adult layers present
	 Tympanic membrane
	 Eyelids fusing laterally and medially
	 Optic tract reaches ventral portion of lateral geniculate body
	 "The hindbrain "presents striking resemblance to that of the newborn."
	 "The rhombencephalon...presents striking resemblance to that of the newborn."
	 Amygdala area
	 Brain represents 43% of embryo
	 Brain: Caudate nucleus and putamen within corpus striatum
	 Cerebellar commissures
	 Cerebellum with external germinal layer
	 Cerebral hemispheres cover lateral portion of diencephalon
	 Choroid plexus now lobular
	 Cortical plate covers nearly all of neopallial surface
	 Dura lines entire vertebral canal
	 Fasciculus cuneatus and fasciculus gracilis form the decussation of the medial lemnisci
	 Greater palatine nerve
	 Grey and white matter
	 Hippocampus reaches temporal pole
	 Inferior and superior cerebellar peduncles
	 Most cisterns present
	 Principal nucleus of inferior olivary nuclei
	 Pyramidal decussations
	 Right- and left-handedness emerges
	 Suprapineal recess
	 Suprascapular nerve
	 Vermis of cerebellum
	 Crown-heel length 4.3 cm
	 Embryo contains approximately 1 billion (10^9) cells
	 Embryonic Period Ends
	 The 8-week embryo has formed more than 4,000 permanent body parts.

Unit 9: 8 to 9 Weeks	
8 weeks, 1 day $\frac{1}{2}$ $\frac{1}{2}$	 Humerus: Bone marrow replaces cartilage
8 weeks, 1 day - 9 weeks $\frac{1}{2}$ $\frac{1}{2}$	 Anal canal patent
8 $\frac{1}{2}$ weeks $\frac{1}{2}$ $\frac{1}{2}$	 Eyelids completely fused
	 Neurons synapse in cerebral cortex (marginal zone)
9 weeks $\frac{1}{2}$ $\frac{1}{2}$	 Bends hip & knee if sole of foot touched
	 Drinking fluid is becoming routine
	 Sucking the thumb
	 The young fetus now sighs, stretches, moves the head, opens the mouth, and moves the tongue
	 Tongue movement
	 Female fetuses have early reproductive cells in their ovaries
	 Thyroid gland weighs 2 grams
	 Small intestine peristalsis
	 External capsule
	 Face, hands, and feet sense light touch
	 Olivary nucleus with five components
Unit 10: 9 to 10 Weeks	
9 weeks - 10 weeks $\frac{1}{2}$ $\frac{1}{2}$	 Early vocal cords
	 Larynx recanalizes
	 My weight will rise more than 75% this week
9 $\frac{1}{2}$ weeks $\frac{1}{2}$ $\frac{1}{2}$	 I yawn when I want
9 weeks, 4 days $\frac{1}{2}$ $\frac{1}{2}$	 Yawns
10 weeks $\frac{1}{2}$ $\frac{1}{2}$	 Eyes roll downward reflexively
	 Palatine tonsils
	 Fingernails and toenails begin to grow!
	 Three-layered epidermis
	 Tiny unique fingerprints have arrived!
	 Now, all the bones are getting harder
	 Tooth buds (secondary teeth)
	 Glomeruli formation begins
	 Physiologic herniation ends

	<div>Commissure of the fornix</div> <div>Corpus callosum begins</div> <div>Crown-heel length 7.5 cm</div>
Unit 11: 10 to 11 Weeks	
10 weeks - 12 weeks $\frac{1}{2}$ $\frac{1}{2}$	<div>Langerhans cells enter epidermis</div>
10 $\frac{1}{2}$ weeks $\frac{1}{2}$ $\frac{1}{2}$	<div>Volar and plantar pads regress</div>
11 weeks $\frac{1}{2}$ $\frac{1}{2}$	<div>The face now makes complex expressions</div>
	<div>Immunological competence</div> <div>Intermediate layer</div> <div>Nose & lips completely formed</div> <div>Now you can tell if your baby is a girl or a boy!</div> <div>Thyroid gland weighs 12 grams</div> <div>Intestines absorb water & glucose</div> <div>Small intestine lined with villi</div> <div>Auditory cells: inner & outer hair cells</div> <div>Crown-heel length</div>
Unit 12: 11 to 12 Weeks	
11 weeks - 12 weeks $\frac{1}{2}$ $\frac{1}{2}$	<div>Weight increases by 60% this week</div>
12 weeks $\frac{1}{2}$ $\frac{1}{2}$	<div>Hands touch the mouth up to 50 times per hour</div>
	<div>T lymphocytes leave thymus</div> <div>Sebaceous glands</div> <div>Many different hormones are present in pituitary gland</div> <div>Thyroid gland produces hormone</div> <div>Palate fuses</div> <div>Upper limbs reach final proportion</div> <div>All facial muscles in final positions</div> <div>Bladder resembles smooth muscle</div> <div>Bowel movements</div> <div>Liver: Bile production begins</div> <div>There are taste buds all over the mouth</div> <div>Corpus callosum</div> <div>Crura cerebri</div> <div>Myelination in spinal cord</div> <div>Crown-heel length 12 cm</div> <div>Head circumference 10 cm</div>
Unit 13: 3 to 4 Months	

13 weeks $\frac{1}{2}$ $\frac{1}{2}$	Teeth are growing
	Cilia lining airways
	Most of body sensitive to touch
	Crown-heel length 15 cm
14 weeks $\frac{1}{2}$ $\frac{1}{2}$	Girls move their jaws more than the boys do
	Light touch to mouth evokes turn toward stimulus
	4-lobed cerebral cortex
	Cerebellum resembles adult structure
	Crown-heel length 17 cm
	Fat deposits in cheeks
15 weeks $\frac{1}{2}$ $\frac{1}{2}$	Stem cells arrive in bone marrow
	Body fat emerges throughout the body
	Glucagon in fetal bloodstream
	Digestive enzymes
	Crown-heel length 19.5 cm
16 weeks $\frac{1}{2}$ $\frac{1}{2}$	Quickening
	Fat deposits upper & lower limbs
	Tooth enamel
	Colon lined with villi
	Bronchial tree nearly complete
	Canalicular stage begins
	Hormonal stress response to invasive procedures
	Crown-heel length 21 cm
Unit 14: 4 to 5 Months	
17 weeks $\frac{1}{2}$ $\frac{1}{2}$	Retina has discrete layers
18 weeks $\frac{1}{2}$ $\frac{1}{2}$	Apocrine sweat glands
	Cream-like substance protects skin
	Sweat glands
	Insulin secretion
	Speaking motion of larynx
	Corpus callosum complete
19 weeks $\frac{1}{2}$ $\frac{1}{2}$	Melanin production
	Number of oogonia peak (at about 7 million) within fetal ovaries

	<div></div> <div>Daily cycles in biological rhythms</div>
	<div></div> <div>Sulci on surface of cerebral hemispheres</div>
20 weeks $\frac{1}{2}$ $\frac{1}{2}$	<div></div> <div>All skin layers and structures</div>
	<div></div> <div>Peyer's patches</div>
	<div></div> <div>Surfactant production (low levels)</div>
	<div></div> <div>Hearing and responding to sound begins</div>
	<div></div> <div>Hearing and responding to sound begins</div>
	<div></div> <div>Crown-heel length 28 cm</div>
	<div></div> <div>Head circumference 20 cm</div>
Unit 15: 5 to 6 Months	
20 weeks - 24 weeks $\frac{1}{2}$ $\frac{1}{2}$	<div></div> <div>Eyelids separate, eyes open and close</div>
21 weeks $\frac{1}{2}$ $\frac{1}{2}$	<div></div> <div>Periderm disappears</div>
	<div></div> <div>Stratum corneum</div>
21 weeks - 22 weeks $\frac{1}{2}$ $\frac{1}{2}$	<div></div> <div>If born prematurely from this point on, survival is possible</div>
22 weeks $\frac{1}{2}$ $\frac{1}{2}$	<div></div> <div>Cornea structure</div>
	<div></div> <div>Behavioral states</div>
23 weeks $\frac{1}{2}$ $\frac{1}{2}$	<div></div> <div>Brain weight 100 grams</div>
24 weeks $\frac{1}{2}$ $\frac{1}{2}$	<div></div> <div>Blink-startle response; females before males</div>
	<div></div> <div>Crown-heel length 34.5 cm</div>
Unit 16: 6 to 7 Months	
25 weeks $\frac{1}{2}$ $\frac{1}{2}$	<div></div> <div>Intestinal lining contains all adult cell types</div>
	<div></div> <div>Rods & cones</div>
	<div></div> <div>The ability to taste</div>
26 weeks $\frac{1}{2}$ $\frac{1}{2}$	<div></div> <div>Additional fat deposits decrease wrinkles</div>
	<div></div> <div>Tear production</div>
	<div></div> <div>Terminal sac stage begins</div>
	<div></div> <div>The ability to smell has arrived</div>
26 weeks - 38 weeks $\frac{1}{2}$ $\frac{1}{2}$	<div></div> <div>Brain weight increases 400% to 500%</div>

27 weeks $\frac{1}{2}$ $\frac{1}{2}$	 Pupils react to light
28 weeks $\frac{1}{2}$ $\frac{1}{2}$	 Distinguishes sounds of different frequencies
	 Crown-heel length 39.5 cm
Unit 17: 7 to 8 Months	
30 weeks $\frac{1}{2}$ $\frac{1}{2}$	 Breathing motions are common even though there is no air in the womb
	 6-layered cerebral cortex
	 Head circumference 30 cm
32 weeks $\frac{1}{2}$ $\frac{1}{2}$	 Esophagus: Lower esophagus muscles functional
	 Glomeruli formation complete
	 Alveoli
	 Memory - music preferences
	 Crown-heel length 45 cm
Unit 18: 8 to 9 Months	
32 weeks - 36 weeks $\frac{1}{2}$ $\frac{1}{2}$	 Prenatal food affects newborn taste preferences
34 weeks $\frac{1}{2}$ $\frac{1}{2}$	 Rapid weight gain
35 weeks $\frac{1}{2}$ $\frac{1}{2}$	 Firm grip
	 Amniotic fluid volume peaks
36 weeks $\frac{1}{2}$ $\frac{1}{2}$	 Surfactant production accelerates
	 Brain weight 300 grams
	 Crown-heel length 48.5 cm
Unit 19: 9 Months to Birth	
37 weeks $\frac{1}{2}$ $\frac{1}{2}$	 Fetus drinks an estimated 15 oz (or 450cc) of amniotic fluid/day
38 weeks $\frac{1}{2}$ $\frac{1}{2}$	 Air breathing begins
	 By term, the typical umbilical cord measures 20 to 24 inches (50 to 60 cm)
	 Heart beats 54 million times before birth
	 Major circulatory changes
	 Spinal cord ends at third lumbar vertebrae
	 Brain weight 350 grams

	<input type="checkbox"/> Crown-heel length 50 cm
	<input type="checkbox"/> Fetus initiates labor
	<input type="checkbox"/> Head circumference 35 cm
	<input type="checkbox"/> Time to be born!
<div> <div>66 weeks, 5 days</div> <div> <div> <div>1/2</div> <div>1/2</div> </div> </div> </div> <div> <input checked="" type="checkbox"/> Premuscle cells form sheets representing muscles of facial expression </div>	